Solution for "geodesic" motion of a Schwarzschild black hole along a magnetic field in $AdS^2 \times \mathbb{S}^2$ space-time

George A. Alekseev

Steklov Mathematical Institute of Russian Academy of Sciences,
Gubkina 8, 119991, Moscow Russia
E-mail: G.A.Alekseev@mi.ras.ru

arXiv:1512.06289v1 [gr-qc] 19 Dec 2015

The exact solution of Einstein - Maxwell equations for a Schwarzschild black hole immersed in the static spatially homogeneous $AdS^2 \times \mathbb{S}^2$ space-time of Bertotti-Robinson magnetic universe is presented. In this solution, the black hole possesses a finite initial boost in the direction of the magnetic field and performs a "geodesic" oscillating motion interacting with the background gravitational and electromagnetic fields.

Keywords: Einstein-Maxwell, Schwarzschild, Bertotti-Robinson universe, rigid frames

Introduction

The century which passed since a wonderful discovery of General Relativity, brought us a lot of developed methods for construction of exact solutions of Einstein and Eisntein-Maxwell field equations and findings of numerous particular solutions.[1–3] However, despite the permanent interest in the literature to different aspects of solutions which describe an interaction of various compact sources (such as black holes, particle-like singularities, matter discs, rings, etc) with the external gravitational and electromagnetic fields, very few such exact solutions are known which describe some non-trivial motion of the sources in various external fields.

In this paper, we present a construction of new solution of Einstein-Maxwell equations for a Schwarzschild black hole moving freely along a magnetic field in $AdS^2 \times \mathbb{S}^2$ space-time of Bertotti-Robinson magnetic universe shown on Fig. 1.



Fig. 1. Schwarzschild black hole in a "geodesic" motion along a magnetic field in Bertotti-Robinson (BR) magnetic universe. In Weyl-like coordinates, the black hole horizon is represented by segments on the axis $\rho = 0$ which constitute a black stripe on (t, z)-plane. $H = g_{\tau\tau}/g^{(BR)}_{\tau\tau}$ and $g_{\tau\tau}$ is the square of time-like Killing vector field $\partial/\partial\mathcal{T}$ which becomes null on the horizon.

Among the earlier found solutions which admit some dynamical interpretation it is worth to mention the Ernst solution in Ref. 4 which describes a charged black hole immersed in Melvin electric universe and accelerated by its electric field (see also Ref. 5 for details and references) and solution derived in Ref. 6 for a Schwarzschild black hole immersed in Bertotti-Robinson magnetic universe and resting at the origin of some rigid frame without any struts or string-like singularities.

In comparison with the solution Ref. 6, the solution presented here possess one more arbitrary real parameter which determines the initial boost of a black hole along the direction of the magnetic field. This boosted black hole performs a free "geodesic" motion, oscillating around the origin of a chosen rigid reference frame. Below, we consider at first the motion of a test particles along the magnetic field in the Bertotti-Robinson background and show that any such particle performs an oscillating motion around the origin of some rigid reference frame where a geodesic particle with zero boost can be at rest. Then we show that a similar rigid frame can be associated with any other geodesic world line directed along the magnetic field. In each of these frames the metric in co-moving coordinates coincides with the same Bertotti-Robinson one. Therefore, we can use this metric as the background and apply the solution generating methods developed earlier for Einstein-Maxwell equations[a] for construction of the solution with a black hole immersed in this metric, that was done in Ref. 6. The generated solution is static in a co-moving frame, but in the original frame it looks like a free oscillating motion of a black hole near its equilibrium position without struts and string-like singularities on the axis.

The background Bertotti-Robinson space-time with $\mathbf{AdS^2 \times \mathbb{S}^2}$ topology and spatially homogeneous magnetic field

The space-time of the Bertotti-Robinson magnetic universe possess $\mathrm{AdS}^2 \times \mathbb{S}^2$ topology and it is filled by spatially homogeneous magnetic field (see Ref. 1,2 for details and references). In cylindrical coordinates $\{t, \rho, z, \varphi\}$, the components of its metric, 1-form of electromagnetic potential and the only non-zero component of magnetic field measured by the local observer can be presented in the form ($\gamma = c = 1$):

$$\begin{aligned} ds^2 &= -\cosh^2\Big(\frac{z}{b}\Big)\, dt^2 + d\rho^2 + dz^2 + b^2 \sin^2\Big(\frac{\rho}{b}\Big)\, d\varphi^2 \\ \underline{\mathbf{A}} &= \Big\{0, 0, 0, -2b \sin^2\Big(\frac{\rho}{2b}\Big)\Big\}, \qquad H_{\hat{z}} = \frac{1}{b} \end{aligned} \tag{1}$$

where b is a constant parameter. The sections $\{t = const, z = const\}$ possess the geometry of a 2-sphere of the radius b and the space-time is closed in the ρ-direction. In the limit $b \to \infty$, the magnetic field vanishes ($H_{\hat{z}} \to 0$) and the metric (1) transforms into the usual Minkowski metric in cylindrical coordinates.

[a]See, in particular, Ref. 7 for generation of Einstein-Maxwell solitons or Ref. 8,9 for application of the monodromy transform approach and the corresponding integral equation method.

Geodesic motion of test particles along the z-axis in the metric (1)

The world lines of test particles which are at rest in the metric (1) constitute a rigid frame. These world lines possess the acceleration along z-axis which value is

$$W^z = \frac{1}{b}\tanh\frac{z}{b}$$

and only the world line of a test particle at rest in the position $z = 0$ is geodesic. The time-like geodesics which correspond to a motion of test particles of mass μ and the energy E_o along z-axis in metric (1) are

$$\tanh\frac{z}{b} = \frac{\sqrt{\mathcal{E}_o^2 - 1}}{\mathcal{E}_o}\sin\frac{t - t_*}{b}, \qquad \mathcal{E}_o = \frac{E_o}{\mu} > 1 \tag{2}$$

This equation shows that the motion of a freely falling neutral test particle boosted with some finite energy $\mathcal{E}_o > 1$ along the z-axis in the metric (1) is periodic and it takes place in some restricted region $\cosh\frac{z}{b} < \mathcal{E}_o$.



Fig. 2. The left picture shows the form of g_{tt} metric component which plays the role of a potential well for test particles moving with the energy $\mathcal{E}_o$ along a magnetic field in Bertotti-Robinson magnetic universe. The right picture shows the geodesic world lines for these test particles on (t, z) plane for different energies $\mathcal{E}_o$.

Killing vector fields in (t, z)-plane

The rigid frames in the metric (1) are determined by congruences of Killing vector fields in (t, z)-plane. In the coordinates (t, ρ, z, φ) these can be expressed as:

$$\xi^i = \{\cosh\delta - \sinh\delta\sin\frac{t - t_*}{b}\tanh\frac{z}{b},\ 0,\ \sinh\delta\cos\frac{t - t_*}{b},\ 0\} \tag{3}$$

where δ and t_* are arbitrary real parameters. Thus, at every point of (t, z)-plane we have a family of Killing vector fields parametrized by δ which possess the norm

$$g_{\xi\xi} \equiv \xi_i\xi^i = -\cosh^2\frac{z}{b}\left(\cosh\delta - \sinh\delta\,\sin\frac{t - t_*}{b}\tanh\frac{z}{b}\right)^2 + \sinh^2\delta\,\cos^2\frac{t - t_*}{b}.$$

Rigid frames associated with geodesics on (t, z)-plane

The background solution (1) possess an important symmetry: changing the reference frame with corresponding coordinate transformation $(t, \rho, z, \varphi) \to (\mathcal{T}, \rho, \mathcal{Z}, \varphi)$ with

$$\begin{cases} \mathcal{T} = b\, \arctan\left[\cosh\delta\ \tan\dfrac{t-t_*}{b} - \sinh\delta\ \tanh\dfrac{z}{b}\ \sec\dfrac{t-t_*}{b}\right], \\ \mathcal{Z} = b \,\mathrm{arcsinh}\left[\cosh\delta\ \sinh\dfrac{z}{b} - \sinh\delta\ \cosh\dfrac{z}{b}\ \sin\dfrac{t-t_*}{b}\right], \end{cases} \tag{4}$$

leavs the form of the background metric invariant:

$$ds^2 = -\cosh^2\frac{z}{b}\ dt^2+dz^2+d\rho^2+\sinh^2\frac{\rho}{b}d\varphi^2 = -\cosh^2\frac{\mathcal{Z}}{b}\ d\mathcal{T}^2+d\mathcal{Z}^2+d\rho^2+\sinh^2\frac{\rho}{b}d\varphi^2$$

Thus, we constructed a rigid frame which motion with respect to the original rigid frame is determined by the boost parameter δ. Among the world lines $\mathcal{Z} = const$ only the line $\mathcal{Z} = 0$ is geodesic and for this we have

$$\mathcal{E}_o = \cosh\delta$$

The other Killing lines with $\mathcal{Z} = const \neq 0$ and with the same δ are not geodesics.

Schwarzschild black hole in a "geodesic" motion in metric (1)

The solution for a Schwarzschild black hole at rest in the external Bertotti-Robinson magnetic universe was constructed in Ref. 6[b]. In a slightly different notations it takes the form

$$ds^2 = -H\left(\cosh\frac{\mathcal{Z}}{b}\right)^2 d\mathcal{T}^2 + f(d\rho^2 + d\mathcal{Z}^2) + \frac{b^2\left(\sin\dfrac{\rho}{b}\right)^2}{H}\, d\varphi^2$$

where the functions H, f and the electromagnetic potential $\mathbf{A} = \{0, 0, 0, A_\varphi\}$ are

$$H = \frac{(x_2+y_1)^2}{x_2^2-1},\ \ A_\varphi = -\frac{b(x_2+1)(1+y_1)}{x_2+y_1},\ \ f = \frac{(x_2+y_1)^2}{x_2^2-y_2^2}\left[\frac{x_1+x_2-y_1-y_2}{x_1+x_2+y_1+y_2}\right]^2$$

and the bipolar coordinates $\{x_1, y_1\}$, $\{x_2, y_2\}$ are functions of ρ and $\mathcal{Z}$ only:

$$\begin{cases} x_1 = -\dfrac{b}{m}\sinh\dfrac{\mathcal{Z}}{b}, \\ y_1 = -\cos\dfrac{\rho}{b}, \end{cases} \qquad \begin{cases} x_2 = (R_1+R_2)/2m, \quad y_2 = (R_1-R_2)/2m \\ R_\pm = \sqrt{(b\sinh\dfrac{\mathcal{Z}}{b}\cos\dfrac{\rho}{b} \mp m)^2 + b^2\cosh^2\dfrac{\mathcal{Z}}{b}\sin^2\dfrac{\rho}{b}} \end{cases}$$

However, this solution will describe the boosted Schwarzschild black hole moving in Bertotti-Robinson background, if $\mathcal{T}$ and $\mathcal{Z}$ are functions of t and z defined in (4). For small mass m, this black hole produces perturbations of metric and electromagnetic field localized near $\mathcal{Z} = 0$ that is a background geodesic oscillating near $z = 0$.

[b]It is necessary to mention here that, in contrast to such "local" (in the region near the axis $\rho = 0$) interpretation of this solution as a black hole in the external field, its global interpretation is more difficult because the semi-closed $\mathrm{AdS}^2 \times \mathbb{S}^2$ topology of this external space-time and focusing properties of gravitational field give rise to existence of naked singularity located on the "antipodal" axis $\rho = \pi b$ (see Ref. 6 for more details).

Acknowledgements

This work was supported in part by the Russian Foundation for Basic Research (Grants No. 14-01-00049 and No. 14-01-00860).

References

1. Stephani H, Kramer D, MacCallum M, Hoenselaers C and Herlt E (2003) *Exact Solutions of Einstein's Field Equations*, 2nd edition (Cambridge University Press).
2. Griffiths J B and Podolsky J (2009) *Exact Space-Times in Einstein's General Relativity* (Cambridge University Press).
3. Alekseev G A, *Thirty years of studies of integrable reductions of Einstein's field equations*, Proceedings of the Twelfth Marcel Grossmann Meeting on General Relativity, edited by Thibault Damour, Robert T Jantzen and Remo Ruffini, World Scientific, Singapore, Part A, Plenary and Review talks, p. 645 - 666, (2011); arXiv:gr-qc/1011.3846.
4. Ernst, F.J. (1976) *Removal of the nodal singularity of the C-metric* JMP **17**, 515.
5. Bicak J and Kofron D (2010) *Rotating charged black holes accelerated by an electric field* Phys.Rev. **D82**, 024006; arXiv: gr-qc/1006.4072v1.
6. Alekseev G A and Garcia A A (1996) *Schwarzschild black hole immersed in a homogeneous electromagnetic field*, Phys.Rev. **D53**, 1853-1867.
7. Alekseev G A (1980) *N-soliton solutions of Einstein - Maxwell equations*, JETP Lett. **32** (4), 277-279.
8. Alekseev G A (1985) *The method of the inverse problem of scattering and the singular integral equations for interacting massless fields*, Sov. Phys. Dokl. **30**, 565.
9. Alekseev G A (1988) *Exact solutions in General Relativity*, Proc. Steklov Inst. Maths. 3, 215-262.